

50X1-HUM

CLASSIFICATION S-E-C-R-E-T **SECRET**
SECURITY INFORMATION
CENTRAL INTELLIGENCE AGENCY
INFORMATION FROM
FOREIGN DOCUMENTS OR RADIO BROADCASTS

REPORT

CD NO. --

COUNTRY German Democratic Republic
SUBJECT Scientific - Chemistry, toxic substances
HOW PUBLISHED Monthly periodical
WHERE PUBLISHED Berlin
DATE PUBLISHED Apr 1951
LANGUAGE German

DATE OF INFORMATION 1951

DATE DIST. 16 Feb 1952

NO. OF PAGES 2

SUPPLEMENT TO REPORT NO.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF ESPIONAGE ACT 50 U. S. C. 31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IS ANY HARMED TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW. REPRODUCTION OF THIS FORM IS PROHIBITED.

THIS IS UNEVALUATED INFORMATION

SOURCE Chemische Technik, Vol III, No 4, 1951, pp 117-22.

NEW EAST GERMAN RODENT POISONS


Dr H. Fuerst
Magdeburg

Recently, new synthetic organic rodent poisons have been replacing the old means for rodent control, such as thallium sulfate, phosphorus, zinc phosphide, sulfur dioxide and hydrogen sulfide, and squill.

"Muritan," made by the Bayer Dye Works at Leverkusen, like AMTU, is a thiourea derivative, i.e., it contains a diazo salt of an aromatic thiourea compound. It is effective at a concentration of only 1.5 mg per kilogram of weight of animal, but in contrast to the stable AMTU it easily undergoes chemical decomposition in the bait. This is an advantage, since bait which has not been taken by the rodents need not be picked up by the exterminator, while this must be done in inorganic poisons are used in the bait.

The "Antiraz" preparation, made by the Wolfen Dye Works, also contain thiourea derivatives.

The Bayer Dye Works have been producing poison in grain form for several years. This preparation contains an organic effective substance called "Gastrix," with the formula



It is 2-chloro-4-dimethyl-amino-6-methyl pyrimidine. Its hydrochloride is used. It is made by condensing thiourea and acetoacetic ester into a pyrimidine ring. An intermediate product is "Uracil," known as a fission product of nucleic acids. Gastrix is more toxic than thallium compounds, but it does not accumulate in the soil.

- 1 -

SECRET

CLASSIFICATION S-E-C-R-E-T

STATE	<input checked="" type="checkbox"/> NAVY	<input checked="" type="checkbox"/> NSRB	DISTRIBUTION																
ARMY	<input checked="" type="checkbox"/> AIR	<input checked="" type="checkbox"/> FBI																	


50X1-HUM

SECRET

S-E-C-R-E-T

W. Maesig, in Anzeiger fuer Schaedlingskunde, Vol 23, 1950, p 49, states that the effective substance contained in the insecticide S 605 is suitable as rat poison, with 10 mg per kilogram weight of animal constituting the lethal dose. However, it is reported that the rats will not eat the bait.

F. Hueter, in Anzeiger fuer Schaedlingskunde, Vol 22, 1949, p 84, cites a report by G. Hecht and H. Henecka in Angewandte Chemie, Vol 61, 1949, p 365, on a highly toxic compound, tetramethylene disulfotetramine. It is made by condensation of sulfamide and formaldehyde, and has the formula



However, the article quoted by Hueter mentions it only as highly toxic, without making specific reference to its use as a rodent poison. Hueter himself states that it is useful as a pest control, and points out that it is highly toxic to higher animals. The manufacture, by the above-mentioned method, is surprisingly easy, but the report gives no details on the method of producing the sulfamide.

- E N D -

- 2 -

S-E-C-R-E-T

SECRET